

# Quantitative Evaluation of Variance in Secondary Dentition Eruption Among Ethnic Groups in Hawai'i

Mark H.K. Greer \*  
Kevin J. Loo \*\*

\*Chief, Dental Health Division, Department of Health, State of Hawai'i, 1700 Lanakila Avenue Room 203 Honolulu, Hawai'i 96817, phone 808 – 832-5700, fax 808 – 832-5722, [mhkgreer@mail.health.state.hi.us](mailto:mhkgreer@mail.health.state.hi.us), \*\*Tufts University School of Dental Medicine, Dental Research Intern

**Abstract:** Though little scientific evidence existed to support the belief among dentists who treat Pacific Islander populations that many children of the region erupt secondary teeth earlier and at an eruption rate which exceeds Caucasian children. Based upon a data set created in Hawai'i during the 1998-1999 school year, of 26,097 public school children, the opportunity presented itself to examine for variance in eruption timing and sequence. Hawai'i is an ethnic diverse community, with a majority population comprised of Asians and Pacific Islanders. Children, 5 through 9 years of age, were examined for gender and ethnic variance. In the aggregate, at all ages, girls erupted teeth earlier than boys, however, while generally true among individual tooth types, that variance was not always statistically significant. By ethnic group, African Americans exhibited earlier eruption by contrast with Caucasians, however, Caucasian children caught up by nine years of age. Native Hawaiian, Samoan and Tongan children exhibited earlier and high rates of secondary dentition eruption than Caucasian or African American children. Children of various Asian cohorts did not exhibit significant variance by contrast with Caucasians. Based upon these findings, the authors recommend that dietary fluoride supplementation of Native Hawaiian, Samoan and Tongan children begin at birth rather than 6 months of age and that these children be targeted for pit & fissure sealants as early as five years of age. (PHD, 2003; 10 (1), Pages 45-50)

## Introduction

Hawai'i holds the distinction of being the only state in the nation where Asians and Pacific Islanders comprise the ethnic majority. Based upon the 2000 United States Census, 51.0 percent of the State population fall into this category. This is a decline from the 61.8 percent reported by the 1990 United States (U.S.) Census. Based upon the 2000 Census, Caucasians, at 24.3 percent of the total resident population represent the largest (unmixed) single ethnic group, followed by Japanese (16.6 percent), Filipinos (14.1 percent), Hispanics (7.2 percent), Chinese (4.6 percent) and Koreans (1.9 percent). Persons of mixed ethnic background account for 21.4 percent of Hawai'i's population. Native Hawaiians (pure and part Hawaiians) and (non-Asian) Pacific Islanders account 6.6 percent and 2.8 percent of Hawai'i's resident population, respectively.<sup>1</sup>

Children in Hawai'i have been shown to suffer from disproportionately high rates of dental caries by contrast with their U.S. mainland counterparts. Standardized oral health surveys have also demonstrated significant variance in caries prevalence, unmet treatment needs and preventive service utilization among Hawai'i's ethnically diverse population.<sup>2, 3, 4</sup> Comprehensive Statewide oral health assessments of public school children have also found higher caries prevalence rates in the primary dentition of boys by contrast with girls and higher caries prevalence rates in the secondary dentition of girls by contrast with boys. This is a typical finding in community caries prevalence studies and is attributable, in part, to variance in eruption timing.<sup>5, 6</sup>

Anecdotal reports have suggested that there is significant variance among Hawai'i's ethnically and culturally diverse population in the sequence and rates of eruption of secondary teeth. A common belief among dental providers in Hawai'i is that children of Pacific Islander ancestry, erupt secondary teeth far earlier and more rapidly than Caucasian and Asian children. Baume noted that contemporary Polynesian children tend to erupt their

permanent molars 1-2 years earlier than their European counterparts and a few reports have been published relating to variance in eruption sequencing and timing among Asian and American populations.<sup>7, 8, 9, 10, 11, 12</sup>

Through this project, standardized data sets were analyzed for a variety of indicators of secondary tooth eruption, including timing, sequencing and quantity over time in search of variance among ethnic groups. The identification of eruption variance among ethnic groups may be used in consideration of updating industry standards and policy associated with the placement of dental sealants and the implementation of dietary fluoride supplementation schedules. In addition, findings may contribute the knowledge base associated with comparative odontology.

## Methods

This cross-sectional study is based upon an analysis of a computerized data set including 26,097 standardized oral health examinations conducted and collected by calibrated public health dental hygienists employed by the Hawai'i State Department of Health during the 1998-1999 school year. The data was analyzed for mean number of all erupted secondary teeth, mean number of erupted secondary teeth by tooth type and the proportion of children with at least one erupted tooth by tooth type; all by age and ethnic group. The data was analyzed for children ages 5, 6, 7, 8 and 9.

A determination of ethnic background was made by a combination of the subjective opinion of the dental hygienist observers and information solicited from children and school officials. With the exception of Native Hawaiian children, which included pure and part Hawaiian children regardless of blood quantum, and Southeast Asians, which collectively included Vietnamese, Laotian, Cambodian, Indonesian, Thai and other children of regional ancestral origin, the remaining ethnic categories were limited to children who were ethnically pure. For the purposes of most of this report, ethnically mixed children were excluded from the data set. That set included 562

African American children, 3,698 Caucasians, 615 Chinese, 486 Hispanics, 4,964 Filipinos, 223 Koreans, 1,536 Japanese, 7,526 Native Hawaiians, 1,060 Samoans, 284 Southeast Asians and 312 Tongans. Overall, the sample included 61 percent males and 39 percent females.

Teeth were identified as erupted when the clinical crown met observational criteria for standardized caries prevalence surveys. These criteria included exposure of the entire occlusal surface of posterior teeth (pre-molars and molars) and at least one-third of the anatomic crown of anterior teeth (incisors and cuspids). The age of children was identified as 'age in years at the child's last birthday'. Data was analyzed contrasting variables by gender and contrasting various ethnic groups against Caucasians as the benchmark for comparison. Variance was considered statistical significance where  $p < 0.05$  and confidence intervals were calculated at 95 percent.

**Results**

As shown in Table 1, at all ages evaluated, girls had erupted a greater mean number of secondary teeth than boys and at all ages, that variance was statistically significant ( $p < 0.001$ ). In this case, data presented include all boys and all girls in the available data set, including mixed race/mixed ethnic group children. Generally,


Hispanic, Chinese, Korean and Japanese ancestry exhibited a mean variance, which varied with no discernible pattern, however, that variance was not statistically significant ( $p > 0.05$ ). Filipino children demonstrated a statistically lower mean number of teeth at 9 years of age. Likewise, Southeast Asian 6 year olds exhibited fewer secondary teeth than Caucasian 6 year olds. Significant variance between African Americans and Caucasians disappeared by 9 years of age. Eruption rates in African American children, which exceeded those found among Caucasians, were significantly lower than those found in Samoan and Tongan children, throughout the age group examined. While the mean rates among Native Hawaiians exceeded those found among African Americans, the difference was not statistically significant until age 9.

By comparison with the other ethnic groups examined, overall, children of Native Hawaiian, Samoan and Tongan ancestry demonstrated significantly higher rates of secondary tooth emergence than other children in Hawai'i. Tables 3, 4 and 5 show rates, which exceed African American, Caucasian and Asian groups. Data among Native Hawaiians, Samoans and Tongans reflect earlier emergence, as evident by higher proportions of children with at least one erupted tooth by tooth-type, shown in Table 6, combined with a higher rates of eruption, shown

**Table 1 Variance Between Boys and Girls in Mean Number of Erupted Secondary Teeth, by Age in Years**

	n	5 Year Olds	6 Year Olds	7 Year Olds	8 Year Olds	9 Year Olds
Boys	2,571	1.11 (1.03, 1.19)	4.79 (4.68, 4.91)	8.82 (8.71, 8.92)	11.63 (11.54, 11.72)	13.87 (13.59, 14.12)
Girls	2,539	1.62 (1.52, 1.71)	5.72 (5.60, 5.85)	9.82 (9.73, 9.92)	12.62 (12.50, 12.73)	15.20 (14.87, 15.53)
		$p < 0.001$	$p < 0.001$	$p < 0.001$	$p < 0.001$	$p < 0.001$
95% Confidence Level						

**Figure 1 Mean Number of Erupted Secondary Teeth Over Time**



variance in mean by tooth-type between girls and boys existed, with girls having more erupted secondary teeth than boys. Such findings are consistent with findings in previous reports.<sup>10, 11</sup>

An examination of the mean number of all erupted secondary teeth by age and ethnic group, shown in Table 2 and Figure 1, shows that African American, Native Hawaiian, Samoan and Tongan children had rates of secondary tooth eruption which exceeded those found in Caucasian children. For ages 5 through 9, children of

in Tables 2, 3, 4 and 5, with mean numbers of erupted secondary teeth which fail to converge in pre-adolescence with other ethnic groups. Depending upon tooth-type, either Samoan or Tongan children account for the highest proportion of children with erupted secondary teeth for each tooth-type presented, shown in Table 6. Fewer Caucasian children appear to have erupted secondary teeth in early childhood by contrast with African American children, however, they appear to catch up to and bypass African American children in later childhood.

**Table 2 Variance by Contrast With Caucasians Benchmark for Mean Number of Erupted Secondary Teeth**

		Hispanic	Chinese	Korean	Japanese				
5 Years of Age	Caucasians	1.08 (± 0.13)	1.36 (± 0.42)	0.94 (± 0.28)	1.05 (± 0.52)	0.99 (± 0.20)			
6 Years of Age	Caucasians	4.81 (± 0.21)	4.75 (± 0.60)	4.32 (± 0.53)	5.44 (± 0.88)	4.52 (± 0.34)			
7 Years of Age	Caucasians	8.83 (± 0.18)	8.39 (± 0.60)	8.60 (± 0.50)	9.26 (± 0.78)	8.78 (± 0.31)			
8 Years of Age	Caucasians	11.58 (± 0.16)	11.48 (± 0.41)	11.37 (± 0.39)	12.17 (± 0.68)	11.77 (± 0.27)			
9 Years of Age	Caucasians	13.09 (± 0.40)	13.30 (± 1.02)	12.84 (± 0.136)	13.29 (± 2.05)	13.82 (± 0.89)			
		African American	Filipino	Southeast Asian	Native Hawaiian	Samoaan	Tongan		
5 Years of Age	Caucasians	1.08 (± 0.13)	1.55 (± 0.40) <i>p</i> < 0.02	1.10 (± 0.12)	1.24 (± 0.59)	1.60 (± 0.13) <i>p</i> < 0.001	2.65 (± 0.38) <i>p</i> < 0.001	2.78 (± 0.77) <i>p</i> < 0.001	
6 Years of Age	Caucasians	4.81 (± 0.21)	5.64 (± 0.56) <i>p</i> < 0.005	4.87 (± 0.19)	3.85 (± 0.81) <i>p</i> < 0.02	5.73 (± 0.16) <i>p</i> < 0.001	7.44 (± 0.38) <i>p</i> < 0.001	7.01 (± 0.75) <i>p</i> < 0.001	
7 Years of Age	Caucasians	8.83 (± 0.18)	9.56 (± 0.49) <i>p</i> < 0.01	8.91 (± 0.16)	8.69 (± 0.57)	9.76 (± 0.12) <i>p</i> < 0.001	10.92 (± 0.33) <i>p</i> < 0.001	11.81 (± 0.68) <i>p</i> < 0.001	
8 Years of Age	Caucasians	11.58 (± 0.16)	12.03 (± 0.53) <i>p</i> < 0.05	11.70 (± 0.15)	11.89 (± 0.58)	12.46 (± 0.13) <i>p</i> < 0.001	14.45 (± 0.46) <i>p</i> < 0.001	15.04 (± 1.01) <i>p</i> < 0.001	
9 Years of Age	Caucasians	13.09 (± 0.40)	13.27 (± 0.87)	14.58 (± 0.51) <i>p</i> < 0.001	14.19 (± 1.29)	15.10 (± 0.39) <i>p</i> < 0.001	17.80 (± 1.35) <i>p</i> < 0.001	17.56 (± 1.78) <i>p</i> < 0.001	

Variance is statistically significant where *p* 0.05  
 Confidence Interval calculated at 95%  
 \* *p* > 0.05

**Table 3 Mean Number of All Erupted Secondary Teeth, Contrast by Gender Among 6 Year Olds**

	African American	Caucasian	Chinese	Hispanic	Filipino	Korean	Japanese	Native Hawaiian	Samoaan	Southeast Asian	Tongan
<b>Mandibular 1st Molars</b>											
Boys	1.24(0.94)	1.20 (0.95)	1.30 (0.89)	1.17(0.92)	1.31 (0.91)	1.44 (0.89)	1.15 (0.94)	1.42 (0.88)	1.71 (0.69)	0.98 (0.96)	1.68 (0.70)
Girls	1.59(0.75)	1.44 (0.87)	1.23 (0.94)	1.42 (0.90)	1.53 (0.81)	1.67 (0.73)	1.34 (0.91)	1.63 (0.75)	1.89 (0.39)	1.00 (0.98)	1.75 (0.61)
	<i>p</i> < 0.02	<i>p</i> < 0.001	*	*	<i>p</i> < 0.001	*	<i>p</i> < 0.05	<i>p</i> < 0.001	<i>p</i> < 0.01	*	*
<b>Maxillary 1st Molars</b>											
Boys	1.00(0.99)	1.11 (0.95)	0.96 (0.95)	1.10 (0.97)	1.08 (0.95)	0.89 (0.97)	0.89 (0.94)	1.28 (0.92)	1.61 (0.75)	0.95 (0.95)	1.47 (0.83)
Girls	1.35(0.90)	1.27 (0.93)	0.85 (0.95)	1.13 (0.97)	1.25 (0.92)	1.62 (0.74)	0.98 (0.97)	1.44 (0.85)	1.67 (0.68)	0.79 (0.95)	1.45 (0.82)
	<i>p</i> < 0.05	<i>p</i> < 0.02	*	*	<i>p</i> < 0.002	<i>p</i> < 0.005	*	<i>p</i> < 0.001	*	*	*
<b>Mandibular Centrals</b>											
Boys	1.59 (0.79)	1.37 (0.90)	1.26 (0.94)	1.38 (0.90)	1.25 (0.94)	1.41 (0.89)	1.26 (0.95)	1.41 (0.89)	1.65 (0.73)	1.19 (0.95)	1.79 (0.58)
Girls	1.78 (0.61)	1.43 (0.88)	1.25 (0.94)	1.40 (0.88)	1.44 (0.87)	1.81 (0.60)	1.38 (0.90)	1.59 (0.79)	1.84 (0.52)	1.15 (0.96)	1.70 (0.70)
	*	*	*	*	<i>p</i> < 0.001	*	*	<i>p</i> < 0.001	<i>p</i> < 0.02	*	*
<b>Maxillary Centrals</b>											
Boys	0.61 (0.89)	0.40 (0.76)	0.32 (0.71)	0.42 (0.81)	0.36 (0.72)	0.44 (0.75)	0.40 (0.77)	0.60 (0.87)	0.88 (0.94)	0.46 (0.81)	0.97 (0.97)
Girls	0.61 (0.89)	0.53 (0.84)	0.51 (0.85)	0.56 (0.90)	0.52 (0.83)	0.47 (0.81)	0.57 (0.85)	0.79 (0.93)	1.11 (0.95)	0.26 (0.67)	0.84 (0.91)
	*	<i>p</i> < 0.02	*	*	<i>p</i> < 0.001	*	*	<i>p</i> < 0.001	*	*	*
<b>Mandibular Laterals</b>											
Boys	0.51 (0.83)	0.33 (0.70)	0.33 (0.67)	0.27 (0.69)	0.38 (0.74)	0.48 (0.80)	0.42 (0.78)	0.48 (0.81)	0.82 (0.96)	0.29 (0.68)	0.74 (0.95)
Girls	0.75 (0.93)	0.44 (0.79)	0.54 (0.83)	0.52 (0.84)	0.56 (0.86)	0.71 (0.90)	0.57 (0.85)	0.67 (0.90)	1.14 (0.93)	0.26 (0.62)	1.04 (0.96)
	*	<i>p</i> < 0.05	*	*	<i>p</i> < 0.001	*	*	<i>p</i> < 0.001	<i>p</i> < 0.01	*	*

Variance is statistically significant where *p* ? 0.05  
 \* *p* > 0.05

**Table 4 Among 6 Year Olds, Variance by Contrast With Caucasian Benchmark in Mean (s.d.) Number of Erupted Teeth, by Tooth Type**

	African American	Chinese	Hispanic	Filipino	Korean	Japanese	Native Hawaiian	Samoan	Southeast Asian	Tongan
Mandibular 1st Molars Caucasians 1.32 (0.92)	1.40 (0.88)	1.27 (0.91)	1.29 (0.91)	1.40 (0.87) <i>p</i> <0.05	1.54 (0.82)	1.24 (0.93)	1.51 (0.83) <i>p</i> <0.001	1.81 (0.55) <i>p</i> <0.001	1.00 (0.97) <i>p</i> <0.005	1.72 (0.65) <i>p</i> <0.001
Maxillary 1st Molars Caucasians 1.19 (0.94)	1.16 (0.97)	0.91 (0.95) <i>p</i> <0.002	1.11 (0.96)	1.16 (0.94)	1.21 (0.94)	0.93 (0.95) <i>p</i> <0.001	1.35 (0.89) <i>p</i> <0.001	1.64 (0.71) <i>p</i> <0.001	0.87 (0.94) <i>p</i> <0.005	1.46 (0.82) <i>p</i> <0.02
Mandibular Centrals Caucasians 1.40 (0.89)	1.68 (0.72) <i>p</i> <0.001	1.26 (0.94)	1.39 (0.89)	1.34 (0.91)	1.58 (0.79)	1.31 (0.92)	1.49 (0.85) <i>p</i> <0.02	1.75 (0.63) <i>p</i> <0.001	1.16 (0.95) <i>p</i> <0.05	1.74 (0.64) <i>p</i> <0.001
Maxillary Centrals Caucasians 0.46 (0.80)	0.61 (0.89) <i>p</i> <0.05	0.39 (0.77)	0.49 (0.85)	0.44 (0.77)	0.46 (0.77)	0.48 (0.81)	0.68 (0.90) <i>p</i> <0.001	1.01 (0.95) <i>p</i> <0.001	0.39 (0.77)	0.90 (0.94) <i>p</i> <0.001
Mandibular Laterals Caucasians 0.38 (0.75)	0.62 (0.88) <i>p</i> <0.001	0.41 (0.74)	0.39 (0.78)	0.46 (0.80) <i>p</i> <0.05	0.58 (0.85)	0.49 (0.82) <i>p</i> <0.02	0.57 (0.86) <i>p</i> <0.001	1.00 (0.95) <i>p</i> <0.001	0.28 (0.64)	0.90 (0.96) <i>p</i> <0.001

\* *p*>0.05  
s.d. standard deviation

**Table 5 Among 8 Year Olds, Variance by Contrast With Caucasian Benchmark in Mean (s.d.) Number of Erupted Teeth, by Tooth Type**

	African American	Chinese	Hispanic	Filipino	Korean	Japanese	Native Hawaiian	Samoan	Southeast Asian	Tongan
Mandibular 1st Molars Caucasians 1.99 (0.10)	2.00 (0.00)	1.98 (0.12)	2.00 (0.00)	1.99 (0.14)	1.98 (0.13)	1.98 (0.22)	1.99 (0.12)	2.00 (0.00)	2.00 (0.00)	2.00 (0.00)
Maxillary 1st Molars Caucasians 1.96 (0.16)	1.98 (0.12)	1.94 (0.32) <i>p</i> <0.01	1.99 (0.09)	1.97 (0.22)	1.93 (0.26) <i>p</i> <0.02	1.97 (0.21)	1.99 (0.14)	2.00 (0.06)	1.99 (0.11)	1.97 (0.24)
Mandibular Centrals Caucasians 1.99 (0.13)	2.00 (0.00)	1.98 (0.19)	2.00 (0.00)	1.99 (0.14)	1.98 (0.13)	1.98 (0.19)	1.99 (0.11)	1.98 (0.12)	2.00 (0.00)	2.00 (0.00)
Maxillary Centrals Caucasians 1.91 (0.38)	1.90 (0.39)	1.848 (0.49) <i>p</i> <0.05	1.84 (0.51)	1.88 (0.43)	1.89 (0.41)	1.89 (0.43)	1.94 (0.32)	1.99 (0.13) <i>p</i> <0.002	1.89 (0.39)	1.97 (0.17)
Maxillary Laterals Caucasians 1.38 (0.88)	1.41 (0.88)	1.31 (0.92)	1.14 (0.88)	1.16 (0.92) <i>p</i> <0.001	1.51 (0.78)	1.40 (0.87)	1.50 (0.82) <i>p</i> <0.001	1.72 (0.82) <i>p</i> <0.001	1.19 (0.95)	1.75 (0.85) <i>p</i> <0.001
Mandibular Laterals Caucasians 1.84 (0.52)	1.90 (0.43)	1.82 (0.56)	1.79 (0.57)	1.84 (0.50)	1.86 (0.48)	1.89 (0.42)	1.89 (0.41) <i>p</i> <0.002	1.98 (0.19) <i>p</i> <0.001	1.87 (0.44)	2.00 (0.00) <i>p</i> <0.001
Maxillary 1st Bicuspid Caucasians 0.16 (0.47)	0.20 (0.56)	0.17 (0.49)	0.21 (0.57)	0.31 (0.64) <i>p</i> <0.001	0.30 (0.63) <i>p</i> <0.05	0.20 (0.55)	0.34 (0.66) <i>p</i> <0.001	0.67 (0.82) <i>p</i> <0.001	0.28 (0.63) <i>p</i> <0.05	0.73 (0.86) <i>p</i> <0.001
Mandibular 1st Bicuspid Caucasians 0.11 (0.40)	0.13 (0.46)	0.12 (0.37)	0.13 (0.43)	0.19 (0.51) <i>p</i> <0.001	0.26 (0.61) <i>p</i> <0.01	0.12 (0.40)	0.27 (0.60) <i>p</i> <0.001	0.66 (0.86) <i>p</i> <0.001	0.20 (0.46)	0.73 (0.87) <i>p</i> <0.001
Maxillary 2nd Bicuspid Caucasians 0.04 (0.22)	0.06 (0.33)	0.015 (0.12)	0.03 (0.21)	0.08 (0.32) <i>p</i> <0.001	0.03 (0.26)	0.04 (0.25)	0.10 (0.38) <i>p</i> <0.001	0.26 (0.59) <i>p</i> <0.001	0.12 (0.40) <i>p</i> <0.005	0.31 (0.67) <i>p</i> <0.001
Mandibular 2nd Bicuspid Caucasians 0.02 (0.18)	0.06 (0.35) <i>p</i> <0.05	0.03 (0.21)	0.01 (0.09)	0.07 (0.30) <i>p</i> <0.001	0.05 (0.22)	0.03 (0.22)	0.09 (0.369) <i>p</i> <0.001	0.36 (0.70) <i>p</i> <0.001	0.11 (0.42) <i>p</i> <0.001	0.40 (0.74) <i>p</i> <0.001
Maxillary Cuspids Caucasians 0.04 (0.23)	0.10 (0.41) <i>p</i> <0.01	0.01 (0.09)	0.01 (0.09)	0.06 (0.28)	0.05 (0.22)	0.06 (0.32)	0.09 (0.37) <i>p</i> <0.001	0.25 (0.82) <i>p</i> <0.001	0.05 (0.28)	0.31 (0.72) <i>p</i> <0.001
Mandibular Cuspids Caucasians 0.13 (0.44)	0.24 (0.64) <i>p</i> <0.02	0.17 (0.50)	0.14 (0.44)	0.16 (0.50)	0.32 (0.66) <i>p</i> <0.005	0.20 (0.57) <i>p</i> <0.02	0.25 (0.62) <i>p</i> <0.001	0.54 (0.84) <i>p</i> <0.001	0.20 (0.59)	0.75 (0.90) <i>p</i> <0.001

\* *p*>0.05  
s.d. standard deviation

Through a review of the proportion of children with at least one erupted secondary tooth by tooth-type between the ages of 5 and 9, a predominant pattern was evident relating to the sequence of emergence, shown in Table 7. Common among the eleven ethnic groups evaluated, was the eruption of the maxillary lateral incisors 6th within the sequence of emergence and the eruption of the mandibular and maxillary second molars 13th and 14th within the sequence, respectively. Caucasians, Filipinos, Native

Hawaiians, Samoans and Tongans, demonstrated a tendency to erupt mandibular 1st molars prior to erupting mandibular central incisors, while in other ethnic groups, mandibular central incisors emerged first. There was also significant variance in the sequence of emergence involving maxillary and mandibular 2nd bicuspid and their correlation with the eruption of maxillary cuspids. Likewise, comparable variance existed between maxillary and mandibular 1st bicuspid as they relate to mandibular

**Table 6 Percentage of Children With At Least One Erupted Secondary Tooth, By Tooth Type**

	Caucasian	African American	Chinese	Hispanic	Filipino	Korean	Japanese	Native Hawaiian	Samoaan	Southeast Asian	Tongan
<b>5 Years of Age</b>											
Mandibular 1st Molars	19.5 %	24.6 %	17.4 %	20.2 %	25.1 %	18.4 %	19.5 %	33.2 %	49.5 %	23.7 %	56.2 %
Maxillary 1st Molars	15.6	18.5	12.3	19.3	16.4	15.8	11.9	36.8	41.3	21.1	32.8
Mandibular Centrals	21.8	33.1	18.7	30.2	17.3	26.3	19.8	27.8	39.4	28.9	39.1
<b>6 Years of Age</b>											
Maxillary Centrals	26.6	33.6	21.7	25.1	25.9	29.2	27.7	38.5	55.4	22.4	51.2
Mandibular Laterals	22.3	35.6	26.3	21.3	26.5	35.4	28.5	32.2	54.6	17.1	35.5
<b>7 Years of Age</b>											
Maxillary Laterals	29.6	38.5	22.0	26.0	23.2	39.4	26.5	39.9	67.3	18.9	61.5
<b>8 Years of Age</b>											
Maxillary 1st Bicuspids	11.2	11.8	11.8	13.3	20.9	21.0	12.9	23.6	44.5	18.7	47.1
Mandibular 1st Bicuspids	8.0	8.7	10.3	9.7	13.5	17.5	8.8	19.1	39.8	17.3	45.6
<b>9 Years of Age</b>											
Maxillary 2nd Bicuspids	20.9	6.7	12.9	11.1	21.4	0.0	8.8	23.0	41.7	19.0	33.3
Mandibular 2nd Bicuspids	8.4	3.3	9.7	3.8	22.6	7.1	11.8	21.7	46.7	14.3	55.6
Maxillary Cuspids	8.8	6.7	6.4	11.1	16.3	14.3	19.1	20.7	40.0	9.5	50.0
Mandibular Cuspids	15.8	20.0	29.0	18.5	34.5	35.7	35.3	39.4	58.3	28.6	66.7
Centrals	central incisors – U.S. tooth numbering 8, 9, 24 & 25 and F.D.I./W.H.O. numbering 11, 21, 31 & 41										
Laterals	lateral incisors – U.S. tooth numbering 7, 10, 23 & 26 and F.D.I./W.H.O. numbering 12, 22, 32 & 42										

**Table 7 Predominant Sequence of Eruption Among Secondary Teeth**

	1st	2nd	3rd	4th	5th	6th	7th	8th	9th	10th	11th	12th	13th	14th
All Girls	3L	8L	3U	8U	7L	7U	5U	5L	6L	6U	4U	4L	2L	2U
All Boys	3L	8L	3U	8U	7L	7U	5U	5L	6L	4U	4L	6U	2L	2U
Caucasian	3L	3U	8L	8U	7L	7U	5U	5L	6L	6U	4L	4U	2L	2U
African American	8L	3L	3U	7L	8U	7U	6L	5U	5L	6U	4U	4L	2L	2U
Chinese	8L	3L	3U	7L	8U	7U	6L	5U	5L	4U	4L	6U	2L	2U
Hispanic	8L	3L	3U	8U	7L	7U	5U	6L	5L	6U	4U	4L	2L	2U
Filipino	3L	8L	3U	7L	8U	7U	5U	5L	6L	4U	6U	4L	2L	2U
Korean	8L	3L	3U	7L	8U	7U	6L	5L	5U	6U	4L	4U	2L	2U
Japanese	8L	3L	3U	8U	7L	7U	5U	6L	5L	6U	4U	4L	2L	2U
Native Hawaiian	3L	8L	3U	8U	7L	7U	5U	5L	6L	4U	4L	6U	2L	2U
Samoaan	3L	8L	3U	7L	8U	7U	5U	5L	6L	4L	4U	6U	2L	2U
Southeast Asian	8L	3L	3U	8U	7L	7U	5U	5L	6L	4U	4L	6U	2L	2U
Tongan	3L	8L	3U	8U	7L	7U	5U	5L	6L	4L	4U	6U	2L	2U
Utilizing a quadrant tooth numbering system, U = maxillary, L = mandibular, 8 = central incisors, 7 = lateral incisors, 6 = cuspids, 5 = 1st bicuspids, 4 = 2nd bicuspids, 3 = 1st molars and 2 = 2nd molars														

cuspids. A clear tendency towards the eruption of mandibular teeth prior to their maxillary counterparts was evident in both gender and all ethnic groups, with the exception of bicuspids, which often exhibited maxillary precedence.

**Discussion**

Findings identify a higher overall rate of secondary (permanent) tooth eruption in girls, at all ages examined, than boys. However, by tooth-type, for many ethnic groups, the variance between girls and boys was statistically insignificant. From a community health planning and policy perspective, higher rates among girls (gender variance) was not demonstrated to be significant enough to warrant special consideration with regards to the implementation and management of disease prevention or dental treatment strategies.

On the other hand, the findings of this project support, in a quantified manner, previous reports of advanced secondary dentition emergence among Pacific Islanders in French Polynesia and speculation that the norms of tooth eruption among other Pacific Islanders groups may also differ from those of European or Asian ancestral cohorts. In general, Native Hawaiian (regardless of blood quantum), Samoaan and Tongan children were shown to exhibit secondary tooth eruption timing which was earlier and eruption rates which were significantly greater than those seen in Hawai'i public school children of European, African or Asian lineage. Early tooth eruption and early tooth development, by contrast with the (Western) Caucasian benchmark, should be a factor considered when planning and implementing dental disease prevention programs targeting children of Native Hawaiian, Samoaan and Tongan heritage.

Since this analysis did not adjust other factors, which might influence secondary tooth eruption, it can not be concluded that ethnic heritage alone is the deciding factor which influences eruption. More research needs to be done in consideration of the influence of anthropomorphic, cultural and environmental influences on the emergence of secondary teeth as well as possible correlations with the sequence and timing of eruption and exfoliation of the primary dentition. However, based upon the consistency of findings among the various indicators analyzed for this project, we feel that ethnic background is a significant factor in, and predictor of, secondary tooth eruption.

Current policy recommendations support dietary fluoride supplementation in communities with low drinking water fluoride levels beginning at 6 months of age.<sup>1</sup> Currently accepted dosage schedules take into consideration a number of factors, including the intake of fluorides from all sources, the relative benefit of fluoride intake, the relative risk of clinically detectable dental fluorosis and known (typically Western/American) norms of tooth development. The relatively high emergence of secondary teeth in some Pacific Islander populations, which correlates with an earlier onset of tooth development and mineralization, warrant consideration for a formal recommendation of dietary fluoride supplementation which is initiated at birth rather than according to the delayed initiation schedule adopted by the American Dental Association and American Academy of Pediatrics in 1994.<sup>2</sup> Likewise, findings support a need for earlier preventive services intervention, including the placement of dental (pit & fissure) sealants on the vulnerable surfaces of permanent posterior teeth, among Native Hawaiian, Samoan and Tongan children as young as five years of age.

### Acknowledgements

The authors are grateful to the staff of the Hawai'i State Department of Health's Dental Hygiene Branch, without whom this project could not have been conducted. Special thanks to Dr. Karen I. Hu and dental hygienists Susan L. Tengan and Joyce T. Takata who trained and calibrated observers and coordinated the creation of the data sets utilized in this project.

### Reference

1. U.S Census Bureau, Census 2000
2. Chung, C, Runck, D, Niswander, *et al.* Genetic and Epidemiologic Studies of Oral Characteristics in Hawai'i's School Children: Caries and Periodontal Disease, *Journal of Dental Research*; 1970 49(6):1374-1385.
3. Greer, M, Tengan, S. Dental Caries in Early Childhood Among Native Hawaiian Children, *Journal of Community Health & Clinical Medicine for the Pacific (Pacific Health Dialog)*; 1998 5(2): 332-336.
4. Greer, M, Louie, R. *Abstract Oral Health Disparities Among Asians & Pacific Islanders in Hawai'i*, presented at U.S. Surgeon General's Conference on Children and Oral Health - The Face of a Child, June 12-13, 2000, Washington, D.C.
5. The Prevalence of Dental Caries in United States Children, The National Dental Caries Prevalence Survey: 1979-1980, National Caries Program, National Institute of Dental Research, *NIH Publication No. 82-2245*, December, 1981.
6. Oral Health of United States Children, The National Survey of Dental Caries in U.S. Children: 1986-1987, Epidemiology and Oral Disease Prevention Program, National Institute of Dental Research, *NIH Publication No. 89-2247*, September, 1989
7. Baume, L. Report on a Dental Survey Among the School Population of French Polynesia, *Archives of Oral Biology*; 1968 13:787-802.
8. Hoffding, J., Mitsunobu, M., Yamaguchi, K. et al Emergence of Permanent Teeth and Onset of Dental Stages in Japanese Children, *Community Dentistry and Oral Epidemiology*; 1984, 2:55-8.
9. Smith, H., Garn, S. Polymorphisms in Eruption Sequence of Permanent Teeth in American Children, *American Journal of Physical Anthropology*, 1987, 74:289-303.
10. Kuthy, R., Ashton, J. Eruption Pattern of Permanent Molars: Implications for School-based Dental Sealant Programs, *Journal of Dental Public Health*; 1989, 49(1): Winter.
11. Smith, J., Brunelle, J. Eruption Patterns of Permanent Teeth in a Population of United States School Children, *Journal of Dental Research*; March (63): 213, Abstract (Special Issue)
12. Garn, S., Sandusky, S. et al Negro-Caucasian Differences in Permanent Tooth Emergence at a Consistent Income Level, *Archives of Oral Biology*; May(18):609-615
13. Riordan, P. Fluoride Supplements in Caries Prevention: A Literature Review and Proposal for a New Dosage Schedule, *Journal of Dental Public Health*; 53(3):174-189
14. American Dental Association, Council on Access, Prevention and Inter-professional Relations; *Caries Diagnosis and Risk Assessment: A Review of Preventive Strategies and Management*, JADA; (1995) 126 (Supplement)

**findings support a need for earlier preventive services intervention, including the placement of dental (pit & fissure) sealants on the vulnerable surfaces of permanent posterior teeth, among Native Hawaiian, Samoan and Tongan children**