

Death from multi-resistant shigellosis in Fiji Islands

CHRIS WATSON*

Abstract

Death from Shigellosis is rare in developed countries, however it causes over a million deaths in developing countries worldwide annually. Death from shigellosis is rare in Fiji.² However, the global problem of emerging multi-drug resistance raises some issues about the management of Shigellosis in this country. Within Fiji, Shigella is a notifiable disease. The Fiji Ministry of Health recorded 68 cases of Shigella in 1996, 173 cases in 1997 and 334 cases in 1998 (no data available for 1999).² There was only one recorded death during this time - in 1998. Resistance to chloramphenicol occurred in 82% of cases. Shigella flexneri in Fiji remains sensitive to cephalothin and cefaclor.

The current antibiotic guidelines in Fiji¹, recommend that antibiotics be used only for cases of moderate and severe dysentery. Shigellosis was suspected soon after presentation however the patient was unable to take oral antibiotics and was treated with intravenous antibiotics (chloramphenicol and ampicillin), which were ineffective due to resistance of the organism. The current antibiotic guidelines for severe dysentery recommend chloramphenicol or nalidixic acid - the later not available in Fiji. However the only intravenous drugs that retain their sensitivity to Shigella - ceftriaxone and cephalothin, are expensive (\$F 45.00 per vial of ceftriaxone) and these are only available in large regional hospitals.

Death from Shigellosis is rare in developed countries, however it causes over a million deaths in developing countries worldwide annually.

*Problem Based Learning Tutor, Fiji School of Medicine, Suva, Fiji. Tel: (+679) 311-700 x 1900. Fax: (+679) 305-781. Email: chris_w@fsm.ac.fj

Introduction

Death from Shigellosis is rare in developed countries, however it causes over a million deaths in developing countries worldwide annually. Sixty nine percent of the deaths occur in children under 5 years of age.¹ Death from shigellosis is rare in Fiji.² However, the global problem of emerging multi-drug resistance raises some issues about the management of Shigellosis in this country. This case highlights some of the chemotherapy-related difficulties encountered during the management of severe shigellosis in Fiji.

Case study

Mr M, a previously fit and well 32 year old male, was admitted to Sigatoka District Hospital with a one day history of sudden onset of diarrhea, fever, chills and rigors. He had passed approximately 10 greenish, watery, bloodstained stools in the 24 hours prior to admission, with no history of abdominal pain or vomiting. There were no known contacts with diarrheal disease.

On presentation he was restless, confused, and hypotensive (BP 70/50mmHg), his temperature was 37.8°C. A provisional diagnosis of dehydration secondary to Shigellosis was made and he was given intravenous (IV) fluids and commenced the antibiotics chloramphenicol and ampicillin each in the dose of 500 mgm IV 6 hourly. Mr M. became profoundly hypotensive and hypoglycaemic several hours' later and required further IV fluids. After 24hrs of hospitalization the patient remained restless and unresponsive to voices with a BP of 100/70.

He was transferred to Lautoka hospital on the second day of admission. Upon admission to Lautoka hospital he was severely dehydrated and in septicemic shock with likely Shigella gastroenteritis. The IV fluids, chloramphenicol, and ampicillin were continued. During the evening of the day of admission to Lautoka hospital, the patient continued to deteriorate developing hypokalaemia. Intravenous ceftriaxone was commenced and the other antibiotics were stopped. He developed Disseminated Intravascular Coagulopathy (DIC) with thrombocytopenia and prolonged

Table 1. Antibiotic resistance of *Shigella* species. Lautoka Hospital, Fiji, 1996-1999.

	Antibiotic resistance	Cases 1996	Cases 1997	Cases 1998	Cases 1999
<i>Shigella flexneri</i>					
Total number of cases		23	90	37	19
	<i>Ampicillin</i>	14 (61%)	85 (94%)	35 (95%)	17 (89%)
	<i>Doxycycline</i>	11 (48%)	75 (83%)	28 (76%)	16 (84%)
	<i>Chloramphenicol</i>	10 (43%)	83 (92%)	32 (86%)	14 (74%)
	<i>Cephalothin</i>	1 (4%)	7 (8%)	1 (3%)	Not tested
	<i>Trimethoprim</i>	6 (26%)	28 (31%)	11 (30%)	Not tested
	<i>Cefaclor</i>	Not tested	Not tested	1 (3%)	1 (5%)
	<i>Nalidixic acid</i>	Not tested	Not tested	1 (3%)	Not tested
<i>Shigella sonnei</i>					
Total number of cases		9	7	8	8
	<i>Ampicillin</i>	0	4 (57%)	1 (13%)	4 (50%)
	<i>Doxycycline</i>	0	2 (29%)	Not tested	Not tested
	<i>Cephalothin</i>	0	3 (43%)	Not tested	Not tested
	<i>Trimethoprim</i>	0	1 (14%)	1 (13%)	Not tested
	<i>Cefaclor</i>	Not tested	Not tested	Not tested	2 (25%)
<i>Shigella dysenteriae</i>					
Total number of cases		1	0	3	0
	<i>Ampicillin</i>	0		2 (67%)	
	<i>Doxycycline</i>	0		2 (67%)	
	<i>Chloramphenicol</i>	0		2 (67%)	
	<i>Trimethoprim</i>	0		2 (67%)	

prothrombin time and this was treated with fresh frozen plasma.

The patient continued to deteriorate and required intubation, ventilation and inotropic support because of persisting hypotension. IV flucloxacillin and metronidazole were added. The patient then developed Acute Respiratory Distress Syndrome (ARDS) and remained persistently hypotensive (BP 71/35) despite inotropic support.

Mr M became asystolic and died on the fourth day after his original admission to Sigatoka hospital. Blood cultures remained negative for pathogens throughout his admissions and stool specimen culture results were only available after the patient had died. Stool culture was positive for *Shigella flexneri*. Sensitivities revealed multi-drug resistance to ampicillin and chloramphenicol although the organism was sensitive to ceftriaxone.

Discussion

Shigella is an aerobic gram-negative bacilli which causes diarrhoeal disease in endemic and epidemic forms in poor and crowded communities. Man is the only known reservoir for infection, which is spread by person-to-person contact from asymptomatic excretors and via contaminated water or food. Four serotypes of *Shigella* are responsible for bacillary dysentery - *Shigella dysenteriae*, *flexneri*, *boydii* and *sonnei*. *Dysenteriae* and *flexneri* are responsible for most infections in the tropics and have a case fatality rate of up to 20%.⁴

Infected patients may be asymptomatic or present with a spectrum of symptoms, from mild watery diarrhoea to severe bloody dysentery with extra-intestinal manifestations. The dysentery is characterized by invasion of the colonic mucosa with local spread of the infecting organism and death of intestinal epithelial cells. Extra-intestinal com-

plications can include seizures, hyponatraemia and hypoglycaemia, septicaemia Reiter's syndrome, encephalopathy and haemolytic uraemic syndrome. An endotoxin is likely to play an important role in the systemic manifestations.

Diagnosis is difficult in areas with poor laboratory facilities, as clinically it is often confused with other bloody diarrheas such as amoebic dysentery or campylobacter [iv]. Diagnosis can only be confirmed by culture and identification from faecal specimens or rectal swabs. There is a facility within Fiji to send specimens from outlying hospitals to one of the government laboratories on the main island though, in the case of *Shigella*, this tends to be only done if a major outbreak is suspected. Stool specimens take approximately two to four days to process. Antibiotics are known to shorten the course of the illness and lower the mortality rate in severe cases, although a knowledge of local sensitivities is particularly important as there is increasing resistance to antibiotics in developing countries.^{5,6,7,8,9} The first multi-drug resistant outbreak occurred in central America from 1969 to 1972 with over 10,000 deaths reported due to strains resistant to chloramphenicol, streptomycin, sulphonomides and tetracycline. Multi-drug resistance amongst all serotypes is an increasing problem worldwide, and there is some evidence that increased case fatality is associated with strains resistant to the antibiotics used initially.¹⁰

Until 1984, ampicillin was the drug of choice for severe dysentery and trimethoprim was used for resistant strains.

Quinolones are currently widely recommended for use in developing countries as they continue to be efficacious, however they are prohibitively expensive and unsuitable for children.⁴ Cephalothin and ceftriaxone remain effective although they are also expensive and no oral preparations are available.

Within Fiji, *Shigella* is a notifiable disease. The Fiji Ministry of Health recorded 68 cases of *Shigella* in 1996, 173 cases in 1997 and 334 cases in 1998 (no data available for 1999).² There was only one recorded death during this time - in 1998. However, the Department of Medicine at Lautoka Hospital has recorded three deaths in young adults who presented with severe dysentery in the past 12 months, although a definitive diagnosis of shigellosis was only made in the case presented above. This may represent an increase in the number of cases or may simply be reflect-underreporting.

A review of the microbiology department records at Lautoka Hospital, (which services approximately 350,000 patients - approximately 40% of the total Fijian population), indicated a total of 169 cases of *Shigella* over the past four

years, and 75% of these were *Shigella flexneri*. There were high levels of multi-drug resistance for all *Shigella* subtypes (see Table 1) and these resistance patterns in Fiji were similar to those found around the world and in other developing nations.^{4,5,7} Resistance to Ampicillin occurred in 89% of cases with some evidence that the rates are increasing over time. Resistance to chloramphenicol occurred in 82% of cases. *Shigella flexneri* in Fiji remains sensitive to cephalothin and cefaclor.

The current antibiotic guidelines in Fiji¹¹, recommend that antibiotics be used only for cases of moderate and severe dysentery. The recommended regimes are: nalidixic acid 1gm 6 hourly given orally, cefaclor SR 375mg 12 hourly given orally, or chloramphenicol 500mg 6 hourly IV although, nalidixic acid is currently unavailable in Government Pharmacies. The guidelines also recommend that sensitivities be obtained because of the problem of drug resistance.

Conclusion

In the case of Mr D, several management problems can be identified. Shigellosis was suspected soon after presentation however the patient was unable to take oral antibiotics and was treated with intravenous antibiotics (chloramphenicol and ampicillin), which were ineffective due to resistance of the organism. The current antibiotic guidelines for severe dysentery recommend chloramphenicol or nalidixic acid - the later not available in

... doctors should be made aware that the drugs available to them in the community will not cover Shigellosis and that if dysentery is severe then immediate transfer is required for proper treatment.

Fiji. These guidelines do not adequately address the problem of emerging multi-drug resistance in Shigellosis, and do not provide clear information on treatment options and of the likely resistance of *Shigella* to Chloramphenicol. However the only intravenous drugs that retain their sensitivity to *Shigella* - ceftriaxone and cephalothin, are expensive (\$F45.00 per vial of ceftriaxone) and these are only available in large regional hospitals. This patient was not transferred to a large hospital for 24 hours despite his critical condition, and appropriate antibiotics were not started for 48 hours after his presentation.

Within Fiji the current statistics indicate that death from Shigellosis is extremely rare. Although there is some suggestion that this may be changing, it does not seem necessary to recommend that more finances be made available to provide intravenous ceftriaxone to the district hospitals. However, doctors should be made aware that the drugs available to them in the community will not cover Shigellosis and that if dysentery is severe then immediate transfer is required for proper treatment.

Acknowledgements

My thanks to Stella Drui for information on antibiotic resistance patterns at Lautoka Hospital, and Dr Michael Lowe for assistance with clinical details.

References

1. Kotloff KL, Winickoff JP, Ivanoff B, Clemens JD, et al. Global burden of Shigella infections: Implications for vaccine development and implementation of control strategies. *Bull World Health Organ* 1999; 77 (8):651-66.
2. Fiji Ministry of Health statistic department. Suva Fiji. *Unpublished data*, June 2000.
3. Murduchowicz G, Huminer D, Siegman-Igra Y, et al. Shigella bacteremia in adults. A report of five cases and review of the literature. *Archives of Internal Medicine*. 1987 Nov;147 (11);2034-7.
4. Shears P, Hart CA. Gastrointestinal bacteria. In Cook GC (ed). *Manson's Tropical Diseases. 20th Edition*. WB Saunders Company Ltd, London, 1997. 835-840
5. Ghosh AR, Sehgal SC. Shigella infections among children in Andaman - an archipeligo of tropical islands in Bay of Bengal. *Epidemiol Infect* 1998 Aug;121(1):43-8.
6. Lim YS, Tay L. Serotype distribution and antimicrobial resistance of Shigella isolates in Singapore. *J Diarrhoeal Dis Res*. 1991 Dec;9(4):328-31.
7. Bogaerts J, Verhaegen J, Munyabikali JP, et al. Antimicrobial resistance and serotypes of Shigella isolates in Kigali, Rwanda (1983-1993): increasing frequency of multiple resistance. *Diagn Microbiol Infect Dis*. 1997 Aug;28 (4).165-71.
8. Taylor DN, Bodhidatta L, Brown JE, et al. Introduction and spread of multi-resistant Shigella dysenteriae in Thailand. *M J Trop Med Hyg*. 1989 Jan; 40(1):77-85.
9. Finkelman Y, Yagupsky P, Fraser D, Dagan R. Epidemiology of Shigella infections in two ethnic groups in a geographic region in southern Israel. *Eur J Clin Microbiol Infect Dis*. 1994 May; 13(5). 367-73.
10. Rawashdeh MO, Abadneh AM, Sherman AA. Shigellosis in Jordanian children: a clinico-epidemiological prospective study and susceptibility to antibiotics. *J Trop Pediatr*. Dec: 40(6): 355-9
11. Fiji Ministry of Health. *Antibiotic Guidelines*. Government Printers, Suva. 1998.

Difficult as it may be to cure, it is always
easy to poison and to kill.

Konrad Adenauer (1876-1967) German statesman